

## Effects of electro-acupuncture on anovulation in women with polycystic ovary syndrome.

[Stener-Victorin E](#)<sup>1</sup>, [Waldenström U](#), [Tägnfors U](#), [Lundeberg T](#), [Lindstedt G](#), [Janson PO](#).

### Author information

- <sup>1</sup>Department of Obstetrics and Gynecology, Göteborg University, Sweden.

### Abstract

#### BACKGROUND:

The present study was designed to evaluate if electro-acupuncture (EA) could affect oligo-/anovulation and related endocrine and neuroendocrine parameters in women with polycystic ovary syndrome (PCOS).

#### METHODS:

Twenty-four women (between the ages of 24 and 40 years) with PCOS and oligo-/amenorrhea were included in this non-randomized, longitudinal, prospective study. The study period was defined as the period extending from 3 months before the first EA treatment, to 3 months after the last EA treatment (10-14 treatments), in total 8-9 months. The menstrual and ovulation patterns were confirmed by recording of vaginal bleedings and by daily registrations of the basal body temperature (BBT). Blood samples were collected within a week before the first EA, within a week after the last EA and 3 months after EA.

#### RESULTS:

Nine women (38%) experienced a good effect. They displayed a mean of 0.66 ovulations/woman and month in the period during and after the EA period compared to a mean of 0.15 before the EA period ( $p=0.004$ ). Before EA, women with a good effect had a significantly lower body-mass index (BMI) ( $p<0.001$ ), waist-to-hip circumference ratio (WHR) ( $p=0.0058$ ), serum testosterone concentration ( $p=0.0098$ ), serum testosterone/sex hormone binding globulin (SHBG) ratio ( $p=0.011$ ) and serum basal insulin concentration ( $p=0.0054$ ), and a significantly higher concentration of serum SHBG ( $p=0.040$ ) than did those women with no effect.

#### CONCLUSION:

Repeated EA treatments induce regular ovulations in more than one third of the women with PCOS. The group of women with good effect had a less androgenic hormonal profile before treatment and a less pronounced metabolic disturbance compared with the group with no effect. For this selected group EA offers an alternative to pharmacological ovulation induction.

PMID:

10716298

[PubMed - indexed for MEDLINE]